arXiv:1302.1336v1 [quant-ph] 6 Feb 2013

Device-independent entanglement quantification

Tobias Moroder,[1] Jean-Daniel Bancal,[2,3] Yeong-Cherng Liang,[2] Martin Hofmann,[1] and Otfried Gühne[1]
[1] *Naturwissenschaftlich-Technische Fakultät, Universität Siegen, Walter-Flex-Str. 3, D-57068 Siegen, Germany*
[2] *Group of Applied Physics, University of Geneva, CH-1211 Geneva, Switzerland*
[3] *Center for Quantum Technologies, National University of Singapore, 3 Science Drive 2, Singapore 117543*

We present a method for quantifying both bipartite and multipartite entanglement in a device-independent manner. Specifically, from the observed correlations or the amount of violation of a given Bell inequality, we present a lower bound on the amount of entanglement present in the underlying system, as measured by the negativity or the genuine negativity. Some of these bounds, such as the one we obtain for the Clauser-Horne-Shimony-Holt Bell inequality, are shown to be tight. These bounds provide further information about the dimension of the underlying Hilbert space or the type of entanglement responsible for the observed correlations. Our method relies on a novel technique to characterize correlations derived from quantum states having additional properties, such as a positive partial transpose.

PACS numbers: 03.67.Mn, 03.65.Ud, 03.67.a

Introduction.—The analysis of most quantum experiments relies on a precise characterization of the equipment employed to generate the data, e.g., see Ref. [1] and references therein. However and maybe quite surprising at first, many tasks in quantum information processing can be achieved without such characterization or knowledge, allowing the conclusions drawn to be device-independent. This approach first started in quantum key distribution where security statements that are independent of the precise implementations are clearly desirable. Indeed, already the security of the Ekert protocol [2] relies on a Bell inequality violation [3] rather than on non-distinguishability of special quantum signals. This direction was further motivated by device testing scenarios where both the true device and the test apparatus come from an untrusted source [4, 5]. In recent years, a variety of other device-independent quantum information tasks has been proposed, including further quantum key distribution [6–9] or self-testing schemes [10, 11], tests for different types of entanglement [12–16], ways to certify the state space dimension [17–19], the nature of measurements [20, 21] as well as randomness expansion [22]. Some of these examples are only partially device-independent but follow the general spirit of removing assumptions about the quantum devices.

In this paper we introduce a technique to characterize the correlations obtained by measuring quantum states which are positive under partial transposition (PPT) [23]. This sets a framework for device-independent entanglement quantification in the bi- and multipartite case. Moreover, this technique provides information on both the dimension and the type of entanglement present in the measured system. Note that such a quantification has so far only been achieved for the simplest experimental scenario together with a qubit assumption [24] or even further measurement properties [25]. We employ the negativity [26] as our primary entanglement measure [27]. This provides further insights into the bipartite Peres conjecture [28], which implies that no PPT entangled state can violate a Bell inequality. Here, we show that a violation by any such state can at best be very small, if not vanishing, for a large class of Bell inequalities.

Problem definition.—Let us start by considering a bipartite Bell-type experiment where each party can employ different measurement settings x, y giving rise to outcomes a, b, respectively; we denote the resulting conditional probability by $P(a, b|x, y)$. This distribution has a quantum representation if there exists a quantum state ρ_{AB} and local measurement operators $M_{a|x}$ and $M_{b|y}$ such that $P(a, b|x, y) = \mathrm{tr}(\rho_{AB} M_{a|x} \otimes M_{b|y})$. Our aim is to characterize the set of $P(a, b|x, y)$ obtainable from ρ_{AB} satisfying certain properties. If ρ_{AB} is only required to be a valid quantum state, we recover the original question leading to Tsirelson's bounds [29–31]. But one can require ρ_{AB} to satisfy more constraints, such as being PPT, i.e., that its partial transpose [23] $\rho_{AB}^{T_A}$ has only non-negative eigenvalues, $\rho_{AB}^{T_A} \geq 0$. One can also ask for a characterization of $P(a, b|x, y)$ if the amount of entanglement is bounded $E(\rho_{AB}) \leq E_{\max}$ using a preferred entanglement measure E. This characterization task generalizes naturally to the multipartite case, e.g., to describe if the tripartite distribution $P(a, b, c|x, y, z)$ is quantum, biseparable [12, 32] or originates from a PPT mixture [33].

Our solution is a superset characterization, analogous to the method of Navascués-Pironio-Acín (NPA) [30]: For instance, in the bipartite case we show that a given distribution $P = P(a, b|x, y)$ can only originate from a PPT state if a special matrix $\chi[P, u]$, that linearly depends on P and on some unknowns u, satisfies $\chi[P, u] \geq 0$ and $\chi[P, u]^{T_A} \geq 0$. If it is impossible to find such parameters u, then P has no PPT quantum representation.

Note that the aimed characterization must hold for all possible dimensions, measurements and states with the desired property. However, since any measurement operator corresponds to a projector in higher dimensions, we can assume the projection property in the following, i.e., the relation $M_{a|x}M_{a'|x} = \delta_{aa'}M_{a|x}$ for the operators $M_{a|x}$ on system A for all x, a and a'. This follows from Naimark's extension [34] that also preserves any entanglement monotone. Also, we shall simultaneously employ the notations $M_{a|x}$ and A_i for measurements operators on system A, likewise for other systems. The set $\{A_i\}$ contains the identity operator $A_0 = \mathbb{1}$ and all but one measurement operator $M_{a|x}$ for each setting. Hence one has the aforementioned projection property and an identity relation $A_iA_0 = A_0A_i = A_i$ for all i.

Technique.—In order to characterize correlations in relation with entanglement we combine the concept of matrix of moments [35–39] with the special partial information of the device-independent setting [30, 31, 40, 41].

Let us start with the matrix of moments for the bipartite case and consider first the scenario where the state ρ_{AB} and measurement operators $M_{a|x}, M_{b|y}$ are known. To this scenario we associate two completely positive (CP) local maps Λ_A, Λ_B that we apply to the quantum state $\chi[\rho] = \chi[\rho_{AB}]_{\bar{A}\bar{B}} = \Lambda_A \otimes \Lambda_B[\rho_{AB}]$. Here $\bar{A}$ and $\bar{B}$ denote the respective output spaces. Specifically, consider the local map $\Lambda_A[\rho] = \sum K_n\,\rho\,K_n^\dagger$ where the Kraus operators are given by $K_n = \sum_i |i\rangle_{\bar{A}\,A}\langle n|\,A_i$, and $|n\rangle_A$, $|i\rangle_{\bar{A}}$ are orthogonal basis states of $\mathcal{H}_A$ and $\mathcal{H}_{\bar{A}}$ respectively. Using a similar map for B one obtains

$$\chi[\rho] = \sum_{ijkl} |ij\rangle_{\bar{A}\bar{B}}\,\langle kl|\,\mathrm{tr}[\rho_{AB}A_k^\dagger A_i \otimes B_l^\dagger B_j]. \qquad (1)$$

Thus the matrix $\chi[\rho]$ is just a matrix of certain expectation values. Since the local maps can also be defined using higher moments, e.g., by choosing Kraus operators $K_n = \sum_{i_1,\ldots,i_\ell} |i_1,\ldots,i_\ell\rangle_{\bar{A}\,A}\langle n|\,A_{i_1}A_{i_2}\ldots A_{i_\ell}$, we shall refer to χ as a moment matrix of level ℓ if it contains all ℓ-fold products of A_i. Since both sets $\{A_i\}, \{B_j\}$ contain the identity, the trace of the underlying state is a matrix entry that we refer to as $\chi[\rho]_{\mathrm{tr}} = \mathrm{tr}[\rho]$. Finally, note that by the structure of these local maps we have a couple of important relations: i) if $\rho \geq 0$ then $\chi[\rho] \geq 0$, ii) if $\rho^{T_A} \geq 0$ then $\chi[\rho]^{T_{\bar{A}}} \geq 0$, and iii) if $\rho^{T_A} \geq 0$ then $\chi[\rho^{T_A}] \geq 0$. This matrix of moment approach can analogously be defined in the multipartite case.

A device-independent characterization draws conclusion only from the observed correlations, hence, many of the entries of χ are unknown a priori. However, even without this information the matrix $\chi[\rho]$ has a structure which follows from known relations that hold independently of state and measurements: 1) A_i, B_j are Hermitian operators, 2) A_i, B_j satisfies the above mentioned projection property and the identity relation, 3) certain entries correspond to the observations $P(a,b|x,y) = \mathrm{tr}[\rho_{AB}M_{a|x} \otimes M_{b|y}]$.

Via this partial information we can decompose without loss of generality each matrix of moments $\chi[\rho]$ as

$$\begin{aligned}\chi[\rho] = \chi[P,u] &= \chi^{\mathrm{fix}}(P) + \chi^{\mathrm{open}}(u)\\ &= \sum_{a,b,x,y} P(a,b|x,y)F_{abxy} + \sum_v u_vF_v,\end{aligned} \qquad (2)$$

i.e., into one fixed part that linearly depends on the observed data $\chi^{\mathrm{fix}}(P) = \sum P(a,b|x,y)F_{abxy}$ and into an orthogonal, open part $\chi^{\mathrm{open}}(u) = \sum_v u_vF_v$ which would be known only by the knowledge of state and measurements. Here all operators $F = F^\dagger$ are Hermitian. Note that the constraint $\chi[\rho]_{\mathrm{tr}} = \chi[P,u]_{\mathrm{tr}} = 1$ is fulfilled automatically if the probabilities P are normalized. We give an example how the relations $1) - 3)$ provide the form given by Eq. (2) in the appendix.

Connection with the NPA hierarchy.—At this point we like to connect the present technique to that of NPA [30, 41], the best known method to characterize quantum correlations. For their method, one can identify a likewise construction $\chi^{\mathrm{NPA}}[\rho] = \Lambda[\rho_{AB}]$, but with Λ being a global CP map which already ensures that if $\rho \geq 0$ then $\chi^{\mathrm{NPA}}[\rho] \geq 0$. If one uses the operator-sum ansatz $\chi^{\mathrm{NPA}}[\rho] = \sum_m L_m\rho\,L_m^\dagger$ where $L_m = \sum_s |s\rangle\,\langle m|\,O_s$ with $|m\rangle, |s\rangle$ being respective basis states for the bipartite in- and output Hilbert spaces, this leads to $\chi^{\mathrm{NPA}}[\rho] = \sum |s\rangle\,\langle t|\,\mathrm{tr}[\rho_{AB}O_t^\dagger O_s]$. If this operator set $\{O_s\}$ consists of all ℓ-fold products of measurement operators, then imposing the constraint $\chi^{\mathrm{NPA}}[\rho] \geq 0$ corresponds to the ℓ^{NPA}-th step in their hierarchy.

Therefore a bipartite moment matrix χ of level ℓ as defined above and a 2ℓ-step χ^{NPA} only differ in the ordering of the expectation values and in that certain moments of χ^{NPA} are not included in χ. Still these similarities are important to relate results about the NPA method χ^{NPA} to the modified moment matrix χ.

PPT Tsirelson bounds & Negativity estimation.—Next we demonstrate how the introduced technique helps to compute Tsirelson bounds, i.e., extremal quantum values of a Bell inequality, or to provide device-independent lower bounds on the amount of entanglement.

The structure of $\chi[\rho] = \chi[P,u]$ as given by Eq. (2) provides a device-independent characterization of quantum correlations: A given probability distribution P can only originate from a quantum state if one can find suitable parameters u such that $\chi[\rho] = \chi[P,u] \geq 0$ holds, otherwise P is non-quantum. Consequently this superset characterization can be used to derive bounds on the optimal quantum value of a Bell inequality, i.e., on a linear function of the observed data, generally parsed as $I \cdot P = \sum I_{abxy}P(a,b|x,y)$, by solving $\max\{I \cdot P \,|\, \chi[\rho] = \chi[P,u] \geq 0\}$. Increasing the level ℓ in $\chi[\rho]$ generates a hierarchy of sharper bounds which, due to the similarity between the $\chi[\rho]$ matrices and the NPA ones, converges to the commutative Tsirelson bound in the limit. Note that this optimization is a semidefinite program which can be solved efficiently [42] .

In comparison with NPA the advantage of the additional structure $\chi = \chi_{\bar{A}\bar{B}}$ is that one can now incorporate further constraints. For instance, one could ask for a similar Tsirelson bound if the underlying state is PPT by including the additional constraint $\chi[\rho]^{T_{\bar{A}}} \geq 0$,

$$\begin{aligned} \max_{P,u} \quad & I \cdot P && (3) \\ \text{s.t.} \quad & \chi[\rho] = \chi[P,u] \geq 0,\ \chi[\rho]^{T_{\bar{A}}} \geq 0,\ \chi[\rho]_{\mathrm{tr}} = 1. \end{aligned}$$

By this method one again obtains an upper bound to the true PPT Tsirelson bound, which converges to the related commutative bound in the limit of large levels ℓ, see appendix for details.

Next, let us show how to estimate the negativity [26], defined via the sum of negative eigenvalues λ_i of the partially transposed state as $N[\rho_{AB}] = \sum_{\lambda_i<0} |\lambda_i(\rho_{AB}^{T_A})|$. In the following we employ its variational form which reads as $N[\rho_{AB}] = \min\{\mathrm{tr}[\sigma_-] | \rho_{AB} = \sigma_+ - \sigma_-, \sigma_\pm^{T_A} \geq 0\}$. Using the properties of the moment matrix, one can readily optimize over a larger set: The constraint $\rho = \sigma_+ - \sigma_-$ is relaxed by $\chi[\rho] = \chi[\sigma_+] - \chi[\sigma_-]$, while $\sigma_\pm^{T_A} \geq 0$ translates to $\chi[\sigma_\pm]^{T_{\bar{A}}} \geq 0$. If one observes a certain violation of a Bell inequality $I \cdot P = v$, a lower bound on the negativity of ρ_{AB} compatible with this observation is given by

$$\begin{aligned} \min_{P,u,P_\pm,u_\pm} \quad & \chi[\sigma_-]_{\mathrm{tr}} && (4) \\ \text{s.t.} \quad & \chi[\rho] = \chi[P,u] = \chi[\sigma_+] - \chi[\sigma_-] \geq 0,\ \chi[\rho]_{\mathrm{tr}} = 1, \\ & \chi[\sigma_\pm]^{T_{\bar{A}}} = \chi[P_\pm, u_\pm]^{T_{\bar{A}}} \geq 0,\ I \cdot P = v. \end{aligned}$$

At this point we like to stress that both optimization problems presented above admit a natural generalization to the multipartite scenario using PPT mixtures (which include biseparable states) and the genuine negativity as a measure for genuine multiparticle entanglement [33]. Further details and the explicit semidefinite programs are given in the appendix.

Examples.—Let us start with the Clauser-Horne-Shimony-Holt (CHSH) inequality [43], where each party has two possible settings $x, y \in \{1, 2\}$ yielding binary outcomes a, b. Using correlation terms $\langle X_x Y_y \rangle = P(a = b|x, y) - P(a \neq b|x, y)$, the inequality reads as

$$I_{\mathrm{CHSH}} = \langle X_1Y_1 \rangle + \langle X_1Y_2 \rangle + \langle X_2Y_1 \rangle - \langle X_2Y_2 \rangle \overset{\mathrm{LHV}}{\leq} 2 \quad (5)$$

for any local hidden-variable model (LHV), while quantum mechanics allows a maximum of $I_{\mathrm{CHSH}}^{\max} = 2\sqrt{2}$. Since every separable state fulfills the LHV bound [44], any violation $I_{\mathrm{CHSH}} > 2$ signals entanglement of the underlying quantum state ρ_{AB}. By solving Eq. (4) we can now provide a quantitative statement in terms of the minimal negativity that the underlying state ρ_{AB} must possess. Specifically, our numerical results lead to the sharp bound $N[\rho_{AB} | I_{\mathrm{CHSH}} = v] \geq (v-2)/(4\sqrt{2}-4)$; the resulting plot and a more detailed discussion, also about the other examples, can be found in the appendix. Note that this recovers the known result that PPT states must necessarily satisfy the CHSH inequality [45].

As a second example, we consider the Bell inequality $I_{3322} \leq 0$ [46, 47] where each party can perform three possible dichotomic measurements as indicated by the subscripts. For $0 \leq v \leq 0.25$, our numerical solution of Eq. (4) gives $N[\rho_{AB} | I_{3322} = v] \geq 2v$. As with the CHSH inequality, we observe that the inequality can only be violated if the underlying state is NPT entangled [precision: 10^{-7}]. The other interesting region is close to the maximum possible quantum value. A Bell state can reach a violation of $I_{3322} = 0.25$ [46], however the best known upper bound is given by $I_{3322}^{\max} \lesssim 0.25088$ and there exist infinite-dimensional states which can asymptotically reach this value [48]. As we see more detailed in the first plot of Fig. 1, if $I_{3322} > 0.25$ the negativity bound satisfies $N[\rho_{AB}] > 1/2$. Because the negativity of any $\mathbb{C}^2 \otimes \mathbb{C}^D$ state is upper bounded by $1/2$ for all $D \geq 2$, this certifies that such violation is achievable only with local Hilbert space dimension $d \geq 3$, or, in other words, that $I_{3322} \leq 0.25$ serves as a dimension witness [17] for qutrits.

This insight inspired us to investigate also the very first Bell inequality used as a dimension witness [17], namely, $I_{2233} \leq 0$ [46, 49, 50]. The corresponding negativity bounds are shown in the second plot of Fig. 1. At a violation of $v = 1/\sqrt{2} - 1/2 \approx 0.2071$ the negativity bound is $1/2$, therefore a larger quantum violation shows that the underlying state is at least 3-dimensional—this also certifies the heuristic qubit bound of I_{2233} presented in Ref. [17]. Also, for the maximal violation at $I_{2233}^{\max} = (\sqrt{11/3} - 1)/3$, our numerical optimization gives a negativity bound that differs from that of the optimal, non-maximally entangled state [51] $\psi_\lambda \propto |00\rangle + \lambda |11\rangle + |22\rangle$ by less than 5×10^{-6}.

Let us point out that the best PPT Tsirelson upper bound that we computed for I_{2233} does not yet coincide with the LHV bound, likewise for a few other Bell inequalities involving a small number of measurement settings. However, for the majority of known facet-defining Bell inequalities involving four dichotomic measurement settings per party [52–54], the respective bounds were found to be vanishing within numerical precision. For details, see Tab. I of the appendix in which we list the best (non-vanishing) values for some of these inequalities.

We also considered examples involving more than two parties, where one is typically interested to verify genuine multipartite entanglement. This strongest form of multiparticle entanglement can be detected from observed correlations by violating a device-independent entanglement witness (DIEW) [12]. By a similar method as in the bipartite case, we investigated the minimal amount of multipartite entanglement (measured by the genuine negativity [33]) needed to violate the DIEWs I_{32} and I_{33}, where each party has respectively two or three dichotomic measurements [12, 55]. Let us point out that I_{32} is the



FIG. 1: Negativity bounds for given violations of Bell inequalities obtained by numerically solving Eq. (4) for different levels of the considered moment matrix. The upper plot shows the I_{3322} inequality [46] for violations close to the maximum, while the lower one corresponds to the I_{2233} inequality [46].

Svetlichny inequality [56], and thus also demonstrates genuine nonlocality. From the bounds we computed (see appendix), we can obtain information about the type of entanglement responsible for given violations, in similar spirit to Ref. [14]. For instance, since the genuine negativity of any state of the three-qubit W-class [57] is bounded by $\sqrt{2}/3$, one verifies that violations close to the maximum of these DIEWs can never be achieved by such type of entanglement. Moreover, our bounds show that these DIEWs can never be violated by states which are PPT mixtures [33]. Using similar arguments as presented in Ref. [15], this result can even be extended to the n-partite witnesses I_{n2} and I_{n3}. This implies that, apart from a quantification, the generalization of PPT Tsirelson bounds to the multipartite case provides a way to approximate the set of biseparable correlations in the presence of more than three parties [12].

Finally, there are also other questions for the multipartite case. At last we computed the maximal violation of the tripartite Bell inequality $I_{S5} \leq 3$ [47] for states which are PPT for all bipartitions. We find that it is bounded by 3.0187, which shows that the example of Ref. [58], optimally violates the tripartite Peres conjecture via this inequality.

Conclusion.—We have presented a versatile tool to characterize quantum correlations in the bi- and multipartite case when the underlying state has additional properties such as a positive partial transpose. The crucial observation is that any matrix of moments, which originates from a quantum state, can be seen as the result of a local map acting on the underlying quantum state if the moments are ordered properly. This key idea finally allows us to provide device-independent bounds on the entanglement present in the measured system. Moreover, the bounds obtained can provide further information about the underlying state space dimension and the type of entanglement involved. Finally, the above observation also facilitates the characterization of biseparable quantum correlations involving arbitrary number of parties.

For future work, we believe that our method can be extended to bound, in a device-independent manner, other entanglement measures. Clearly, it will also be interesting to investigate how our technique can be used in conjunction with other separability criteria, or applied in the closely-related steering scenarios [59].

This work greatly profited from discussions with R. Augusiak and J. I. de Vicente at the Centre of Sciences in Benasque. We would also like to thank M. Navascués, K. F. Pál, V. B. Scholz and T. Vértesi for stimulating discussions about topic and technicalities. This work has been supported by the EU (Marie Curie CIG 293993/ENFOQI), the BMBF (Chist-Era project QUASAR), the Chist-Era project DIQIP, the Swiss NCCR-QSIT, the National Research Foundation and the Ministry of Education of Singapore.

[1] D. Rosset, R. Ferretti-Schöbitz, J.-D. Bancal, N. Gisin, and Y.-C. Liang, Phys. Rev. A **86**, 062325 (2012).
[2] A. K. Ekert, Phys. Rev. Lett. **67**, 661 (1991).
[3] J. S. Bell, Physics **1**, 195 (1964).
[4] C. Mayers and A. Yao, Quantum Inf. Comput. **4**, 273 (2004).
[5] F. Magniez, D. Mayers, M. Mosca, and H. Ollivier, arXiv:quant-ph/0512111.
[6] A. Acín, N. Gisin, and L. Masanes, Phys. Rev. Lett. **97**, 120405 (2006).
[7] S. Pironio, A. Acín, N. Brunner, N. Gisin, S. Massar, and V. Scarani, New J. Phys. **11**, 045021 (2009).
[8] M. Pawłowski and N. Brunner, Phys. Rev. A **84**, 010302(R) (2011).
[9] H.-K. Lo, M. Curty, and B. Qi, Phys. Rev. Lett. **108**, 130503 (2012).

[10] M. McKague, T. H. Yang, and V. Scarani, J. Phys. A: Math. Theor. **45**, 455304 (2012).
[11] T. H. Yang and M. Navascués, arXiv:1210.4409.
[12] J.-D. Bancal, N. Gisin, Y.-C. Liang, and S. Pironio, Phys. Rev. Lett. **106**, 250404 (2011).
[13] K. F. Pál and T. Vértesi, Phys. Rev. A **83**, 062123 (2011).
[14] N. Brunner, J. Sharam, and T. Vértesi, Phys. Rev. Lett. **108**, 110501 (2012).
[15] J.-D. Bancal, C. Branciard, N. Brunner, N. Gisin, and Y.-C. Liang, J. Phys. A: Math. Theor. **45**, 125301 (2012).
[16] C. Branciard, D. Rosset, Y.-C. Liang, and N. Gisin, arXiv:1110.5512 (2012).
[17] N. Brunner, S. Pironio, A. Acín, N. Nicolas Gisin, A. A. Méthot, and V. Scarani, Phys. Rev. Lett. **100**, 210503 (2008).
[18] S. Wehner, M. Christandl, and A. C. Doherty, Phys. Rev. A **78**, 062112 (2008).
[19] R. Gallego, N. Brunner, C. Hadley, and A. Acín, Phys. Rev. Lett. **105**, 230501 (2010).
[20] T. Vértesi and M. Navascués, Phys. Rev. A **83**, 062112 (2011).
[21] R. Rabelo, M. Ho, D. Cavalcanti, N. Brunner, and V. Scarani, Phys. Rev. Lett. **107**, 050502 (2011).
[22] S. Pironio, A. Acín, S. S. Massar, A. Boyer de la Giroday, D. N. Matsukevich, P. Maunz, S. Olmschenk, D. Hayes, L. Luo, T. A. Manning, et al., Nature **464**, 1021 (2010).
[23] A. Peres, Phys. Rev. Lett. **77**, 1413 (1996).
[24] Y.-C. Liang, T. Vértesi, and N. Brunner, Phys. Rev. A **83**, 022108 (2011).
[25] P. Lougovski and S. J. van Enk, Phys. Rev. A **80**, 034302 (2009).
[26] G. Vidal and R. F. Werner, Phys. Rev. A **65**, 032314 (2002).
[27] M. B. Plenio and S. Virmani, Quant. Inf. Comput. **7**, 1 (2007).
[28] A. Peres, Found. of Phys. **29**, 589 (1999).
[29] B. S. Cirel'son, Lett. Mat. Phys. **4**, 93 (1980).
[30] M. Navascués, S. Pironio, and A. Acín, Phys. Rev. Lett. **98**, 010401 (2007).
[31] A. C. Doherty, Y.-C. Liang, B. Toner, and S. Wehner, in *Proc. IEEE Int. Conf. on Computational Complexity* (2008), pp. 199–210.
[32] W. Dür, J. I. Cirac, and R. Tarrach, Phys. Rev. Lett. **83**, 3562 (1999).
[33] B. Jungnitsch, T. Moroder, and O. Gühne, Phys. Rev. Lett. **106**, 190502 (2011).
[34] A. Peres, Found. of Phys. **20**, 1441 (1990).
[35] E. Shchukin and W. Vogel, Phys. Rev. Lett. **95**, 230502 (2005).
[36] J. Rigas, O. Gühne, and N. Lütkenhaus, Phys. Rev. A **73**, 012341 (2006).
[37] H. Häseler, T. Moroder, and N. Lütkenhaus, Phys. Rev. A **77**, 032303 (2008).
[38] A. Miranowicz, M. Piani, P. Horodecki, and R. Horodecki, Phys. Rev. A **80**, 052303 (2009).
[39] N. Killoran and N. Lütkenhaus, Phys. Rev. A **83**, 052320 (2011).
[40] S. Wehner, Phys. Rev. A **73**, 022110 (2006).
[41] M. Navascués, S. Pironio, and A. Acín, New J. Phys. **10**, 073013 (2008).
[42] S. Boyd and S. Vandenberghe, *Convex optimization* (Cambridge University Press, 2004).
[43] J. F. Clauser, M. A. Horne, A. Shimony, and R. A. Holt, Phys. Rev. Lett. **23**, 880 (1969).
[44] R. F. Werner, Phys. Rev. A **40**, 4277 (1989).
[45] R. F. Werner and M. M. Wolf, Phys. Rev. A **61**, 062102 (2000).
[46] D. Collins and N. Gisin, J. Phys. A: Math. Gen. **37**, 1775 (2004).
[47] C. Śliwa, Phys. Lett. A **317**, 165 (2003).
[48] K. F. Pál and T. Vértesi, Phys. Rev. A **82**, 022116 (2010).
[49] D. Kaszlikowski, L. C. Kwek, J.-L. Chen, M. Żukowski, and C. H. Oh, Phys. Rev. A **65**, 032118 (2002).
[50] D. Collins, N. Gisin, N. Linden, S. Massar, and S. Popescu, Phys. Rev. Lett. **88**, 040404 (2002).
[51] A. Acín, T. Durt, N. Gisin, and J. I. Latorre, Phys. Rev. A **65**, 052325 (2002).
[52] N. Brunner and N. Gisin, Phys. Lett. A **372**, 3162 (2008).
[53] K. F. Pál and T. Vértesi, Phys. Rev. A **79**, 022120 (2009).
[54] T. Vértesi (private communication).
[55] J. T. Barreiro, J.-D. Bancal, et al., in preparation.
[56] G. Svetlichny, Phys. Rev. D **35**, 3066 (1987).
[57] A. Acín, D. Bruß, M. Lewenstein, and A. Sanpera, Phys. Rev. Lett. **87**, 040401 (2001).
[58] T. Vértesi and N. Brunner, Phys. Rev. Lett. **108**, 030403 (2012).
[59] H. M. Wiseman, S. J. Jones, and A. C. Doherty, Phys. Rev. Lett. **98**, 140402 (2007).
[60] M. McKague, M. Mosca, and N. Gisin, Phys. Rev. Lett. **102**, 020505 (2009).
[61] K. Horodecki, M. Horodecki, and P. Horodecki, Quant. Inf. Comput. **10**, 0901 (2010).
[62] J.-D. Bancal, N. Gisin, and S. Pironio, J. Phys. A: Math. Theor. **43**, 385303 (2010).
[63] V. B. Scholz and R. F. Werner, arXiv:0812.4305.
[64] M. Navascués, T. Cooney, D. Pérez-García, and N. Villanueva, Found. of Phys. **42**, 985 (2012).
[65] M. Junge, M. Navascues, C. Palazuelos, D. Perez-Garcia, V. B. Scholz, and R. F. Werner, J. Math. Phys. **52**, 012102 (2011).

Appendix

Examples of constraints on the matrix of moments

As an example of how the relations 1) – 3) from the main text lead to the general structure of $\chi[\rho]$ as given by Eq. (2), we consider the case where the map Λ_A is applied to a single system ρ_A, i.e., $\chi[\rho] = \sum_{ij} |i\rangle\langle j| \operatorname{tr}[\rho_A A_j^\dagger A_i]$. Suppose that the first setting x has three outcomes $\{0,1,2\}$ whereas the second one, here labeled as x', has only two outcomes $\{0,1\}$. Using $\{A_i\} = \{\mathbb{1}, M_{0|x}, M_{1|x}, M_{0|x'}\}$ one gets

$$\chi[\rho] = \begin{bmatrix} \mathbb{1}^\dagger \mathbb{1} & \mathbb{1}^\dagger M_{0|x} & \mathbb{1}^\dagger M_{1|x} & \mathbb{1}^\dagger M_{0|x'} \\ M_{0|x}^\dagger \mathbb{1} & M_{0|x}^\dagger M_{0|x} & M_{0|x}^\dagger M_{1|x} & M_{0|x}^\dagger M_{0|x'} \\ M_{1|x}^\dagger \mathbb{1} & M_{1|x}^\dagger M_{0|x} & M_{1|x}^\dagger M_{1|x} & M_{1|x}^\dagger M_{0|x'} \\ M_{0|x'}^\dagger \mathbb{1} & M_{0|x'}^\dagger M_{0|x} & M_{0|x'}^\dagger M_{1|x} & M_{0|x'}^\dagger M_{0|x'} \end{bmatrix}_{\rho_A} \quad (6)$$

where ρ_A indicates that we still must take the expectation values. Via the listed properties 1) − 3) one obtains

$$\chi[\rho] = \begin{bmatrix} \mathrm{tr}(\rho) & P(0|x) & P(1|x) & P(0|x') \\ P(0|x) & P(0|x) & 0 & u_1 + \mathbb{i}u_2 \\ P(1|x) & 0 & P(1|x) & u_3 + \mathbb{i}u_4 \\ P(0|x') & u_1 - \mathbb{i}u_2 & u_3 - \mathbb{i}u_4 & P(0|x') \end{bmatrix}. \quad (7)$$

For the diagonal entries one employs $\mathrm{tr}[\rho_A M_{0|x} M_{0|x}] = \mathrm{tr}[\rho_A M_{0|x}] = P(0|x)$, while the zero entries occur by the projection identity $M_{0|x} M_{1|x} = 0$. Since the expectation value of $M_{0|x} M_{0|x'}$ is not directly accessible we can only set it equal to a general complex entry $u_1 + \mathbb{i}u_2$ using real coefficients u.

Note that whenever the underlying state ρ is normalized, the first term of $\chi[\rho]$ is fixed to be $\mathrm{tr}(\rho) = 1$. In the case where the underlying operator does not need to be normalized, however, as it happens for instance with $\sigma_\pm$ in the negativity estimation by Eq. (3), this entry is not fixed a priori. It is thus given by an unknown variable $u_0 \in \mathbb{R}$.

The multipartite scenario

In this section we define more precisely moment matrices for an arbitrary number of parties, and present multipartite optimization problems that compute the Tsirelson bounds for PPT mixtures and the device-independent quantification of entanglement in terms of genuine negativity.

In a multipartite scenario one can ask, in analogy with Eq. (3), for a bound on the observed correlations if the underlying state is a PPT mixture [33]. A n-partite state ρ is said to be a PPT mixture if it can be written as a convex combination $\rho = \sum_m p_m \rho_m$ of states ρ_m which are PPT with respect to different bipartitions $m \subset \{1, \dots, n\}$ of the n subsystems. Since any state that is separable with respect to a chosen bipartition m is also PPT according to this splitting, the set of biseparable states is included in the set of PPT mixtures. Hence if one verifies that a given state is not a PPT mixture, one automatically certifies that it is not biseparable, and thus, by definition, genuine multipartite entangled.

The definition of moment matrices as given in the main text naturally extend to this n-party scenario by applying a local CP map Λ_s to each subsystem $s = 1, \dots, n$, i.e., $\rho \mapsto \chi[\rho] = \otimes_{s=1}^n \Lambda_s[\rho]$. By the local structure of this transformation one obtains: For any PPT mixture ρ the resulting matrix of moments can be decomposed as $\chi[\rho] = \sum_m p_m \chi[\rho_m]$ with $\chi[\rho_m] \geq 0$ and $\chi[\rho_m]^{T_{\bar{m}}} \geq 0$, with $\bar{m}$ referring to the bipartition on the output spaces. Similar properties as given by 1) − 3) in the main text, constrain the general structure of $\chi[\rho]$ to be $\chi[P, u]$ and one readily obtains a superset approximation for correlations that can be attained via PPT mixtures. Thus if one is interested in the optimal values of a linear expression like $I \cdot P$, where P is generated by a PPT mixture, one obtains a bound by solving

$$\begin{aligned} \max \quad & I \cdot P & (8) \\ \text{s.t.,} \quad & \chi[\rho] = \chi[P, u] = \sum \chi[p_m \rho_m] = \sum \chi[P_m, u_m], \\ & \chi[\rho]_{\mathrm{tr}} = 1,\ \chi[p_m \rho_m] \geq 0,\ \chi[p_m \rho_m]^{T_{\bar{m}}} \geq 0 \quad \forall m. \end{aligned}$$

Note that in this formulation we included already the probabilities p_m into the matrix $\chi[P_m, u_m]$ such that $\chi[P_m, u_m]_{\mathrm{tr}} = p_m$ in this case. We show later in the appendix that even this multipartite extension converges in the limit of an infinite number of moments in $\chi[\rho]$.

For the multipartite equivalent of Eq. (4), let us first remind that the genuine negativity [33] is a computable measure of genuine multipartite entanglement which reduces to the negativity in the bipartite case. It is given by $N_G[\rho] = \min_{\{p_m, \rho_m\}} \sum_m p_m N_m[\rho_m]$ where N_m denotes the negativity with respect to bipartition m and the minimization runs over all possible valid state decompositions of the density operator $\rho = \sum_m p_m \rho_m$ [1]. In analogy to Eq. (4), if one observes a value of $I \cdot P = v$, a lower bound on the genuine negativity compatible with this violation is given by

$$\begin{aligned} \min \quad & \sum \chi[\sigma_m^-]_{\mathrm{tr}} & (9) \\ \text{s.t.} \quad & \chi[\rho] = \sum \chi[p_m \rho_m],\ I \cdot P = v,\ \chi[\rho]_{\mathrm{tr}} = 1, \\ & \chi[p_m \rho_m] = \chi[\sigma_m^+] - \chi[\sigma_m^-] \geq 0,\ \chi[\sigma_m^\pm]^{T_{\bar{m}}} \geq 0 \ \forall m. \end{aligned}$$

Further examples and additional information on the bounds of negativity/ Bell-inequality violation

In this part we present more details on the negativity bounds mentioned in the main text together with some further examples of PPT Tsirelson bounds.

First note that any explicit quantum state ρ_0 which attains a certain value v of a Bell inequality $I \cdot P = v$ provides an upper bound on the minimal negativity compatible with this violation, i.e., $N_{\min}[\rho \,|\, I \cdot P = v] = \min N[\rho \,|\, I \cdot P = v] \leq N[\rho_0]$. Furthermore, since the negativity is convex and invariant by adding local auxiliary states, the minimal negativity is a convex function in the amount of violation v, that we shall denote by $f(v) = N_{\min}[\rho \,|\, I \cdot P = v]$. If one uses the moment matrices $\chi[\rho]$ with increasing levels ℓ one obtains lower bounds

---

[1] The dual of the given measure, i.e., the way how it is defined in Ref. [33], would be $N_G(\rho) = -\min\{\mathrm{tr}[\rho W] | \forall m : W = P_m + Q_m^{T_m}, \mathbb{1} \geq Q_m \geq 0, P_m \geq 0\}$. Hence one sees a missing constraint $\mathbb{1} \geq P_m$ as compared to its original definition, but this does not alter the properties of this measure, i.e., zero for biseparable states, full LOCC monotone, convexity and invariance under local unitaries.

$f_\ell(v) \leq f(v)$ with increasing accuracy. However, if one finds in the bound $f_\ell(v)$ an interval $v \in [a,b]$ such that $f_\ell(v)$ is linear and where the endpoints $v = a, v = b$ are known to be attainable by explicit quantum states then $f_\ell(v) = f(v)$ is a tight bound of the minimal negativity for $v \in [a,b]$.

As we see in more detail in Fig. 2, for both, the CHSH inequality and the I_{3322} inequality for a value between $[0, 1/4]$, the negativity bounds that we obtained by solving Eq. (4) correspond to straight lines for the level $\ell = 3$. Specifically, for the CHSH inequality, this negativity bound is a straight line joining the points $(I_{\text{chsh}} = 2, N[\rho] = 0)$ and $(I_{\text{chsh}} = 2\sqrt{2}, N[\rho] = 1/2)$ with largest numerical deviation $\approx 7 \times 10^{-7}$ among all the computed instances. Likewise, for the I_{3322} inequality, it is a linear bound connecting the origin and the coordinate $(I_{3322} = 1/4, N[\rho] = 1/2)$ with largest numerical deviation $\approx 3 \times 10^{-4}$. Since both endpoints $I_{\text{CHSH}} = 2\sqrt{2}$ and $I_{3322} = 1/4$ can be achieved with a maximally entangled two-qubit state, having negativity $N[\rho] = 1/2$, we thus arrive at the sharp negativity bounds presented in the main text.



FIG. 2: Negativity bounds for CHSH and the I_{3322} Bell inequality as given by Eq. (4) for different levels of the moment matrix considered.

The amount of genuine negativity necessary to achieve different violations of the I_{32} and I_{33} DIEWs were computed according to Eq. (9). For biseparable states these expressions are bounded by $I_{32} \leq 4, I_{33} \leq 6\sqrt{3}$, while the maximal quantum values are $I_{32}^{\max} = 4\sqrt{2}, I_{33}^{\max} = 9\sqrt{3}$ respectively [12, 55, 56]. The results for different hierarchy levels are shown in Fig. 3. Here we refer to partial levels such as level 2+ to denote moment matrices χ which were constructed with all terms involving 2-fold products of local measurement operators, as well as some 3-fold ones. Similarly to the CHSH and I_{3322} inequalities, the plot for the Svetlichny inequality I_{32} at level $\ell = 3$ is a straight line up to numerical precision. Since the value $I_{32} = 4\sqrt{2}$ is achievable with a three-qubit GHZ state with genuine negativity $N_G[\rho] = 1/2$, we obtain the following tight bound for the minimal genuine negativity compatible with a Svetlichny inequality violation, $N_G[\rho_{ABC}|I_{32} = v] \geq (v-4)/(8\sqrt{2}-8)$.

From Fig. 3 we also note that no violation of these DIEW is possible with PPT mixtures; hence the bound for biseparable states coincides with the one for PPT mixtures. Since any DIEW with this property can be found by using Eq. (8), and this hierarchy applies to an arbitrary number of parties, this means that our technique can be used to find DIEWs of this kind for an arbitrary number of parties, in contrast to the numerical three-party technique presented in Ref. [12].

Furthermore, as pointed out in the main text, since any three qubit state of the W-class [32] satisfies [2] $N_G(\rho_W) \leq \sqrt{2}/3 \approx 0.471$, our result shows that violations $v > 5.563$ and $v > 15.36$ of the I_{32} and I_{33} inequalities can never be obtained by any such three-qubit states. Therefore one gains even further information from the achieved bound about the underlying type of entanglement, in similar spirit to Ref. [14].

Finally, note that apart from the bipartite Bell inequalities mentioned in the main text, we have also computed, via Eq. (3), upper bounds on PPT Tsirelson bounds for 175 facet-defining Bell inequalities involving four dichotomic measurement settings per party. These inequalities were taken from Refs. [52–54]. Interestingly, only three of these inequalities give a PPT Tsirelson upper bound which differs from the respective LHV bound by more than 10^{-6}, whereas the numerical precision of the computation is of the order of $10^{-7} - 10^{-10}$. In Tab. I we summarize the best bound that we have computed for

---

[2] Since N_G is convex, this optimization can be performed over pure three-qubit states $|\psi_W\rangle$ of the W-class. Note that the genuine negativity of a pure state is just the minimum bipartite negativity for all bipartitions m. Because N_G is furthermore invariant under local basis changes one can employ the LU-equivalent standard form of a pure three-qubit state of the W-class [57], which leaves a straightforward optimization. The bound is saturated by the W-state.



FIG. 3: Genuine negativity as a function of the observed violation of the tripartite Svetlichny I_{32} and DIEW I_{33} inequalities. Since any three-qubit state of the W-class has a genuine negativity $\lesssim 0.471$, larger violations certify that the underlying state is not of this SLOCC class.

these three inequalities as well as for other Bell inequalities mentioned in the main text. This table also includes the respective Tsirelson bound for the tripartite setting of I_{S5} with respect to states that are PPT for all bipartitions. This optimization problem is like Eq. (3) with the tripartite moment matrix and having a PPT constraint $\chi[\rho]^{T_m} \geq 0$ for each bipartition.

Via this numerical investigation we hoped to find a counterexample to the bipartite Peres conjecture; a PPT state which could violate a Bell inequality. This perspective was triggered by the NPA hierarchy (or using also the modified moment matrix) to compute standard Tsirelson bounds. Although this method is only guaranteed to be complete in the limit of an infinite number of moments, it is important to stress that there are many known instances where one could stop the hierarchy already earlier, since one has already reached the true Tsirelson bound. This is certified by a special rank property of the solution [41] and means that the bound does not improve further even if one includes higher moments. However, in all our considered non-trivial PPT examples the respective bounds sharpened if we considered higher levels.

Bell inequality	PPT Tsirelson bound	ℓ	Matrix size
$I_{\text{chsh}} \leq 2$ [43]	2	1	$3^2 \times 3^2$
$I_{3322} \leq 0$ [46]	0	2	$10^2 \times 10^2$
$I_{2233} \leq 0$ [46]	$\leq 1.3559 \times 10^{-3}$	3+	$45^2 \times 45^2$
$A_6 \leq 0$ [52]	$\leq 7.6754 \times 10^{-6}$	2+	$31^2 \times 31^2$
$I^1_{4422} \leq 0$ [52]	$\leq 9.7947 \times 10^{-6}$	2+	$29^2 \times 29^2$
$I^3_{4422} \leq 0$ [52]	$\leq 2.8531 \times 10^{-4}$	2+	$27^2 \times 27^2$
$I_{S5} \leq 3$ [47]	≤ 3.0187	3	$7^3 \times 7^3$

TABLE I: Upper bounds on the maximal possible violation by PPT states for different bipartite Bell inequalities via the method given by Eq. (3). Here matrix size refers to the dimension of the moment matrix $\chi[\rho]$ involved in the computation whereas ℓ stands for the corresponding level of the moment matrix (e.g., level 2+ means that it contains all 2-fold products of local measurement operators, but also some terms which only appear when considering 3-fold products of local measurement operators). The last inequality corresponds to the tripartite case which is PPT for all bipartitions. The precision of the computation is of the order of 10^{-7} – 10^{-10}.

Statement of real states and measurements

In this section we show that the underlying quantum state and measurement operators can generally be assumed to be real when considering the device-independent quantification of entanglement in terms of (genuine) negativity, i.e., there exists an equivalent real construction having the same (or less) amount of (genuine) negativity. This extends the result of Ref. [60] which already proves that probability distributions arising from quantum theory in a Bell-type experiment can always be reproduced using only real states and real measurement operators.

This real property helps in the numerical implementation of the bi- or multipartite programs, Eqs. (3),(4) and Eqs. (8), (9) respectively, since it provides a notable parameter reduction in the optimization problems. This reduction becomes even greater in the presence of additional symmetries. We shortly comment on this parameter reduction at the end of this section.

Proposition 1. *Any n-partite probability distribution P having a quantum representation with respect to density matrix ρ and measurement operators $A^1_i, A^2_j, \ldots, A^n_k$ (with the superscript labeling the party), also has a real quantum representation, i.e., a representation in terms of a real-valued quantum state $\widehat{\rho} = \widehat{\rho}^T$ and real-valued measurement operators $\widehat{A}^1_i = \widehat{A}^{1T}_i, \widehat{A}^2_j = \widehat{A}^{2T}_j, \ldots, \widehat{A}^n_k = \widehat{A}^{nT}_k$*

having an equal or lower amount of (genuine) negativity, $N_m[\widehat{\rho}] = N_m[\rho]$ for any bipartition m and $N_G[\widehat{\rho}] \leq N_G[\rho]$. Furthermore one has:

a) If ρ is PPT across a bipartition m then $\widehat{\rho}$ can even be assumed to be PPT invariant across m, i.e., $\widehat{\rho} = \widehat{\rho}^{T_m} = \widehat{\rho}^T$. Likewise, if ρ is a PPT mixture then $\widehat{\rho}$ can even be assumed to have a mixture of real PPT invariant states, $\widehat{\rho} = \sum p_m \widehat{\rho}_m$ with $\widehat{\rho}_m = \widehat{\rho}_m^{T_m} = \widehat{\rho}_m^T$ for all m.

b) If the observed distribution is invariant under arbitrary exchange of the parties, $P = V(\pi) P V(\pi)^\dagger$ for all possible permutation π, then the underlying state and measurements can further be assumed to be permutationaly invariant, i.e., $\widehat{V}(\pi)\,\widehat{\rho}\,\widehat{V}(\pi)^\dagger = \widehat{\rho}$ for all π and $\widehat{A}_i^1 = \widehat{A}_i^2 = \ldots = \widehat{A}_i^n$.

Proof. For the first statement we construct for any set of projectors A^s, $s = 1, \ldots, n$ and any state ρ, another real state $\widehat{\rho} = \widehat{\rho}^T$ and real projectors $\widehat{A}^s = \widehat{A}^{sT}$ such that $\mathrm{tr}[\rho A^1 \otimes A^2 \otimes \ldots \otimes A^n] = \mathrm{tr}[\widehat{\rho} \widehat{A}^1 \otimes \widehat{A}^2 \otimes \ldots \otimes \widehat{A}^n]$ with $N_m[\widehat{\rho}] = N_m[\rho]$ for any bipartition m and $N_G[\widehat{\rho}] \leq N_G[\rho]$.

Let us start with the projectors. For any chosen basis we can decompose the matrix $A = A_r + iA_i$ into a real, symmetric part $A_r = \mathbb{Re}(A) = A_r^T$ and an imaginary, anti-Hermitian part $A_i = \mathbb{Im}(A) = -A_i^T$. To this matrix we now associate the real and symmetric matrix

$$\widehat{A} = \mathbb{1} \otimes A_r + Y \otimes A_i = \begin{bmatrix} A_r & A_i \\ -A_i & A_r \end{bmatrix} = \widehat{A}^T \tag{10}$$

where $Y = i\sigma_y$. This represents the well-known isomorphism between Hermitian and symmetric matrices, e.g., Ref. [42]. Note that the operator $\widehat{A}$ acts on the enlarged Hilbert space $\mathcal{H}_{\widehat{A}} = \mathcal{H}_{A'} \otimes \mathcal{H}_A$ with $\mathcal{H}_{A'} = \mathbb{C}^2$. Since the original A satisfies the projection identity $A^2 = A$ this provides the relations $A_r^2 - A_i^2 = A_r$ and $A_r A_i + A_i A_r = A_i$ and henceforth the projection identity for $\widehat{A}$ by

$$\begin{aligned} \widehat{A}^2 &= (\mathbb{1} \otimes A_r + Y \otimes A_i)(\mathbb{1} \otimes A_r + Y \otimes A_i) \\ &= \mathbb{1} \otimes (A_r^2 - A_i^2) + Y \otimes (A_r A_i + A_r A_i) = \widehat{A}. \end{aligned} \tag{11}$$

We employ this construction for all projectors $A^s \mapsto \widehat{A}^s$.

Next let us define the appropriate extension of the density matrix ρ to $\widehat{\rho} = \widehat{\rho}_{A_1' A_1 A_2' A_2 \ldots A_n' A_n}$ where the subscripts label the party to which the Hilbert space belongs. However, for ease of reading, we shall use a different ordering of the Hilbert spaces (namely, the primed ones followed by the unprimed ones) $\widehat{\rho}_{A_1' A_2' \ldots A_n' A_1 A_2 \ldots A_n} = \mathcal{R}(\rho)$, which could be undone $\widehat{\rho} = V \widehat{\rho}_{A_1' \ldots A_n' A_1 \ldots A_n} V$ by an appropriate permutation V on Hilbert spaces. More precisely, this operator is given by

$$\widehat{\rho}_{A_1' A_2' \ldots A_n' A_1 A_2 \ldots A_n} = 2^{-n}\, \mathbb{Re} \left[(\mathbb{1} - iY)^{\otimes n} \otimes \rho \right]. \tag{12}$$

Let us first show that $\widehat{\rho}$ is positive semi-definite and has trace one as required for any density operator. To this end, first note that the right-hand-side of Eq. (12) can be rewritten as a sum of two orthogonal parts, namely, $2^{-n-1}(\mathbb{1} - iY)^{\otimes n} \otimes \rho$ and its transpose, where the orthogonality follows from the fact that $(\mathbb{1} - iY)(\mathbb{1} + iY) = 0$. Since the set of eigenvalues remains unchanged under transposition, this means that the eigenvalues of $\widehat{\rho}$ are precisely one-half of the respective eigenvalues of $2^{-n}(\mathbb{1} - iY)^{\otimes n} \otimes \rho$, but with twice the multiplicity. However since $(\mathbb{1} - iY)/2$ is just a rank-one projector one has that the non-vanishing eigenvalues of $2^{-n}(\mathbb{1} - iY)^{\otimes n} \otimes K$ for any Hermitian matrix K are precisely those of K. Thus, we obtain $\widehat{\rho} \geq 0$, $\|\widehat{\rho}\|_{\mathrm{tr}} = \|\rho\|_{\mathrm{tr}} = 1$ and $\|\mathcal{R}(\rho^{T_m})\|_{\mathrm{tr}} = \|\rho^{T_m}\|_{\mathrm{tr}}$ for any bipartition m.

Next we need to show that these choices indeed preserve the expectation values. This is best seen using the state given by Eq. (12) and the measurement form of Eq. (10) together with the following identity

$$\begin{aligned} &\mathrm{tr}_{A'} \left[(\mathbb{1} - iY) \otimes \rho \; (\mathbb{1} \otimes A_r + Y \otimes A_i) \right] \\ &\qquad = \mathrm{tr}_{A'} [\mathbb{1}]\, \rho\, A_r + \mathrm{tr}_{A'} [-iYY]\, \rho\, A_i \\ &\qquad\quad + \mathrm{tr}_{A'} [Y]\, \rho\, A_i + \mathrm{tr}_{A'} [-iY]\, \rho\, A_r \\ &\qquad = 2\rho\, A, \end{aligned} \tag{13}$$

which follows from $Y^2 = -\mathbb{1}$ and $\mathrm{tr}(Y) = 0$. Applying such identities when tracing out each auxiliary space yields

$$\begin{aligned} \mathrm{tr}[\widehat{\rho} \widehat{A}^1 \otimes \ldots \otimes \widehat{A}^n] =& 2^{-n}\, \mathbb{Re} \left\{ \mathrm{tr} \left[V (\mathbb{1} - iY)^{\otimes n} \otimes \rho V \times \right.\right. \\ & \left.\left. (\mathbb{1} \otimes A_r^1 + Y \otimes A_i^1) \otimes \ldots \right] \right\} \\ =& \mathbb{Re} \left(\mathrm{tr}[\rho\ A^1 \otimes \ldots \otimes A^n] \right) \\ =& \mathrm{tr}[\rho\ A^1 \otimes \ldots \otimes A^n]. \end{aligned} \tag{14}$$

Next, we prove that the negativity remains constant $N_m[\rho] = N_m[\widehat{\rho}]$ for any bipartition m. Thus we need the partial transposition of $\widehat{\rho}$ with respect to m. For simplicity, we now consider partial transposition with respect to $\widehat{A}$; the general treatment is completely analogous. To this end, we remind that the partial transposition is a linear operation and thus from Eq. (12) and the fact that $Y^T = -Y$, we have

$$\begin{aligned} \widehat{\rho}_{A_1' \ldots A_n' A_1 \ldots A_n}^{T_{A_1' A_1}} &= 2^{-n}\, \mathbb{Re} \left[(\mathbb{1} + iY) \otimes (\mathbb{1} - iY)^{\otimes n-1} \otimes \rho \right], \\ &= U\, \mathcal{R}[\rho^{T_{A_1}}]\, U^\dagger \end{aligned} \tag{15}$$

where $U = \sigma_x \otimes \mathbb{1}^{n-1} \otimes \mathbb{1}_{A_1 \ldots A_n}$ is a unitary matrix and we have made use of the fact that $\sigma_x\, Y\, \sigma_x = -Y$. Since a unitary does not change the eigenvalues and using the previously mentioned invariance of the trace-norm we get $\|\widehat{\rho}^{T_{\widehat{A}}}\|_{\mathrm{tr}} = \|\mathcal{R}(\rho^{T_A})\|_{\mathrm{tr}} = \|\rho^{T_A}\|_{\mathrm{tr}}$, and hence the statement that the negativity is unchanged. This statement holds for an arbitrary bipartition m. Finally, suppose that $\rho = \sum p_m \rho_m$ is the optimal decomposition for the original state ρ in the definition of the genuine negativity, i.e., $N_G[\rho] = \sum p_m N_m[\rho_m]$. Note that because the map $\rho \mapsto \widehat{\rho}$ is linear and preserves positivity one knows that

$\widehat{\rho} = \sum_m p_m \widehat{\rho}_m$ is a valid decomposition of $\widehat{\rho}$ into states attributed to different bipartitions m. Since the genuine negativity is defined by the minimum over all such decompositions, and because $N_m[\widehat{\rho}_m] = N_m[\rho_m]$ one readily obtains $N_G[\widehat{\rho}] \leq \sum p_m N_m[\widehat{\rho}_m] = N[\rho]$, which proves the first part of the proposition.

Henceforth, we shall work with the real state $\widehat{\rho}$ and the corresponding real measurement operators $\widehat{A}_i^s = \widehat{A}_i^{sT}$, $s = 1, \ldots, n$ that can produce the same correlations. Assuming this, statement a) is obtained as follows: If the original state ρ is PPT across a bipartition m, then the real state $\widehat{\rho}$ has to be PPT as well. Therefore $\widehat{\rho}^{T_m} \geq 0$ can be considered as another quantum state, which, similarly would produce the correct observations since all measurements are real,

$$\begin{aligned} P &= \mathrm{tr}[\widehat{\rho} \widehat{A}_i^1 \otimes \widehat{A}_j^2 \otimes \ldots \otimes \widehat{A}_k^n] \\ &= \mathrm{tr}[\widehat{\rho} (\widehat{A}_i^1 \otimes \widehat{A}_j^2 \otimes \ldots \otimes \widehat{A}_k^n)^{T_m}] \\ &= \mathrm{tr}[\widehat{\rho}^{T_m} \widehat{A}_i^1 \otimes \widehat{A}_j^2 \otimes \ldots \otimes \widehat{A}_k^n]. \end{aligned} \tag{16}$$

Here, between the first and the second line we transposed the measurement operator of all parties belonging to the bipartition m. Thus also the equal-weight mixture $\widehat{\rho}^{\mathrm{ave}} = (\widehat{\rho} + \widehat{\rho}^{T_m})/2$ is a real state that is PPT invariant with respect to m and that yields the same correlations. A similar statement holds for PPT mixtures with partial transposition applied to each state ρ_m in the decomposition.

Statement b) follows from a similar argument. For simplicity, we provide the proof below only for $n = 2$ parties; the generalization to an arbitrary number of parties is analogous. Let us suppose that $\widehat{\rho} = \widehat{\rho}_{AB}$ and that this state as well as $\widehat{A}_i, \widehat{B}_j$ are not yet symmetric under exchange of the parties.[3] Let us add appropriate local auxiliary states $|0\rangle, |1\rangle$ (on system A', B' respectively) to signal whether the state is swapped or not, and consider the convex combination $\widehat{\rho}^{\mathrm{sym}}_{\widehat{A}\widehat{B}}$ given by

$$\begin{aligned} \widehat{\rho}^{\mathrm{sym}}_{\widehat{A}\widehat{B}} &= \frac{1}{2} \left[\widehat{\rho}_{AB} \otimes |01\rangle_{A'B'}\langle 01| + V \widehat{\rho}_{AB} V^\dagger \otimes |10\rangle_{A'B'}\langle 10| \right] \\ &= \frac{1}{2} [\widehat{\rho}_{AB} \otimes |01\rangle_{A'B'}\langle 01| \\ &\qquad + \widehat{V} (\widehat{\rho}_{AB} \otimes |01\rangle_{A'B'}\langle 01|) \widehat{V}^\dagger] \end{aligned} \tag{17}$$

where $\widehat{V} = V(\pi_{\widehat{A}\widehat{B}})$ now denotes the swap operator on $\widehat{A} = AA'$ and $\widehat{B} = BB'$. Similarly for the measurements

$$\widehat{A}_i^{\mathrm{sym}} = \widehat{B}_i^{\mathrm{sym}} = \widehat{A}_i \otimes |0\rangle\langle 0| + \widehat{B}_i \otimes |1\rangle\langle 1|, \tag{18}$$

such that one finally can check that this is a valid quantum representation with

$$P = [P + VPV^\dagger]/2 = \mathrm{tr}[\widehat{\rho}^{\mathrm{sym}}_{AB} \widehat{A}_i^{\mathrm{sym}} \otimes \widehat{A}_j^{\mathrm{sym}}]. \tag{19}$$

From the structure given by Eq. (17) one observes that $\widehat{\rho}^{\mathrm{sym}}_{\widehat{A}\widehat{B}}$ is indeed invariant under the swap operator $\widehat{V}$. Finally since the negativity is convex and symmetric [61] one obtains $N[\widehat{\rho}^{\mathrm{sym}}_{\widehat{A}\widehat{B}}] \leq N[\widehat{\rho}_{AB}]$, while the LOCC monotonicity by measuring the primed systems first provides $N[\widehat{\rho}^{\mathrm{sym}}_{\widehat{A}\widehat{B}}] \geq N[\widehat{\rho}_{AB}]$ such that the negativity is indeed invariant. □

Let us briefly remark how Proposition 1 reduces the number of free parameters for the semidefinite programs.

Firstly, note that since the underlying state and both measurements can be chosen to be real, the matrix of moments can be assumed to be symmetric $\chi[\rho] = \chi[\rho]^T$. For the multipartite PPT mixture question of Eqs. (8),(9), this property can be applied to every bipartition, i.e., $\chi[\rho_m] = \chi[\rho_m]^T$. Concerning the negativity finally, this property can be further imposed for the operators appearing in the variational formulation, since a solution given by $\rho_m = \sigma_m^+ - \sigma_m^-$ with $(\sigma_m^\pm)^{T_m} \geq 0$ would imply the alternative solution $\rho_m = \rho_m^T = (\sigma_m^+)^T - (\sigma_m^-)^T$ with $[(\sigma_m^\pm)^T]^{T_m} = [(\sigma_m^\pm)^{T_m}]^T \geq 0$, and hence also its equal-weight mixture. Therefore in the negativity estimation as given by the semidefinite programs of Eqs. (4),(9), we can set additionally that $\chi[\rho_m] = \chi[\rho_m]^T$ and $\chi[\sigma_m^\pm] = \chi[\sigma_m^\pm]^T$.

The statement a) also simplifies the computation of the respective PPT Tsirelson bounds as given by Eq. (3) or Eq. (8). Since the state can be assumed to be real and PPT invariant, this provides the symmetry $\chi[\rho_m] = \chi[\rho_m]^T = \chi[\rho_m]^{T_{\bar{m}}} \geq 0$ for all m.

Finally, whenever one considers a symmetric Bell inequality [62], i.e., satisfying $I_{abxy} = I_{bayx}$ in the bipartite case, one can impose this symmetry also for the corresponding distribution $P(a, b|x, y) = P(b, a|y, x)$, such that b) of Proposition 1 gets relevant. If both local maps in the construction of $\chi[\rho]$ have an equal number of moments, then $\chi[\rho]$ can be assumed to be invariant $V_{\bar{A}\bar{B}} \chi[\rho] V_{\bar{A}\bar{B}}^\dagger = \chi[\rho]$ under the swap operator $V_{\bar{A}\bar{B}}$. This symmetry in particular helps in the higher levels of the hierarchy.

Convergence of respective PPT characterizations

As emphasized in the main text, the method to approximate PPT Tsirelson bounds via Eq. (3) converges to a description of the commutative set in the limit of an infinite number of moments. A similar statement holds for the multipartite case and PPT mixtures. To show this, we start this section by formulating the commutative version of the respective quantum representation. We then proceed to show the equivalence between this commutative and the tensor product version in the case of a finite-dimensional quantum representation and, at last, prove the convergence. We stress that this convergence follows from the convergence of the NPA hierarchy [41].

---

[3] At this point we can already assume that the local Hilbert spaces are isomorphic $\mathcal{H}_A \cong \mathcal{H}_B$ by an appropriate embedding of a possibly smaller space in higher dimension. In this way the swap operator $V = V(\pi_{AB})$ is well defined.

At first let us motivate this distinction between the tensor product and commutative version: In quantum information we are commonly used to employ tensor products for measurements on different subsystems, i.e., $\mathrm{tr}[\rho_{AB} A \otimes B]$. In contrast, in algebraic field theory for instance, measurements on different parts are described by commuting operators $[\tilde{A}, \tilde{B}] = 0$, both acting already on the bipartite state $\tilde{\rho}$ (therefore we use the tilde) such that expectation values become $\mathrm{tr}[\tilde{\rho}\tilde{A}\tilde{B}]$. If one identifies $\tilde{A} = A \otimes \mathbb{1}$, $\tilde{B} = \mathbb{1} \otimes B$ and $\tilde{\rho} = \rho_{AB}$ then one sees that all expectation values using tensor products can be recovered with commuting observables. The converse question, however, is still open and is known as Tsirelson's problem [63–65].

For clarity, we start with the translation of the bipartite PPT constraint $\rho^{T_A} \geq 0$ in commutative terms. We employ the simplified notions $A_i, B_j, \ldots$ introduced in the main text for measurement operators, and $P = \mathrm{tr}[\ldots A_i \ldots B_j \ldots]$ to indicate the functional form of the observed probability distribution.

Definition 1. *i) A bipartite probability distribution P has a* tensor product PPT quantum representation *if there exists a real state ρ_{AB}, local real measurements operators A_i, B_j such that $P = \mathrm{tr}[\rho_{AB} A_i \otimes B_j]$ and that $\rho_{AB} = \rho_{AB}^{T_A}$ is PPT invariant.*

ii) A bipartite distribution P has a commutative PPT quantum representation *if there exists a real state $\tilde{\rho}$, commuting real measurement operators $\tilde{A}_i, \tilde{B}_j$, $[\tilde{A}_i, \tilde{B}_j] = 0$, such that $P = \mathrm{tr}[\tilde{\rho}\tilde{A}_i\tilde{B}_j]$ and for all possible products, indexed by $i = (i_1, i_2, \ldots, i_n)$, $\tilde{A}(i) = \tilde{A}_{i_1}\tilde{A}_{i_2}\ldots\tilde{A}_{i_n}$, and similarly for $\tilde{B}$, it holds*

$$\mathrm{tr}[\tilde{\rho}\tilde{A}(i)\tilde{B}(j)] = \mathrm{tr}[\tilde{\rho}\tilde{A}(i^T)\tilde{B}(j)] \tag{20}$$

with $i^T = (i_n, i_{n-1}, \ldots, i_1)$.

Proposition 2. *Any bipartite probability distribution which has a tensor product PPT quantum representation also has a commutative PPT quantum representation. The converse holds at least if the commutative PPT quantum representation is finite-dimensional.*

Proof. Let us start with the direction i) to ii). The formulation i) is already simplified using Proposition 1. If we now interpret $\tilde{A} = A \otimes \mathbb{1}$, $\tilde{B} = \mathbb{1} \otimes B$ and $\tilde{\rho} = \rho_{AB}$ then these operators naturally fulfil the requirements of being real, that their commutator vanishes and that they provide the correct expectation values. If one employs the PPT invariance $\rho_{AB} = \rho_{AB}^{T_A}$ and $A_i = A_i^T$ one obtains the condition given by Eq. (20) via

$$\begin{aligned}
\mathrm{tr}[\tilde{\rho}\tilde{A}(i)\tilde{B}(j)] &= \mathrm{tr}[\rho_{AB} A(i) \otimes B(j)] = \mathrm{tr}[\rho_{AB}^{T_A} A(i) \otimes B(j)] \\
&= \mathrm{tr}[\rho_{AB} (A_{i_1} A_{i_2} \ldots A_{i_n})^T \otimes B(j)] \\
&= \mathrm{tr}[\rho_{AB} (A_{i_n} A_{i_{n-1}} \ldots A_{i_1}) \otimes B(j)] \\
&= \mathrm{tr}[\rho_{AB} A(i^T) \otimes B(j)] \\
&= \mathrm{tr}[\tilde{\rho}\tilde{A}(i^T)\tilde{B}(j)].
\end{aligned} \tag{21}$$

This finishes the proof in the direction i) to ii).

Let us now turn to the converse. Thus suppose that $\tilde{\rho}$ and $\tilde{A}_i, \tilde{B}_j$, all acting on the finite-dimensional Hilbert space $\tilde{\mathcal{H}}$, are given as in ii). From these operators we now construct local measurement operators and a bipartite quantum state which is PPT; the statement that they even have the extra symmetries follows from point a) of Proposition 1. This construction is very analogous to the corresponding one of the standard Tsirelson problem. However, in order to show that the conditions given by Eq. (20) finally prove that the constructed bipartite state is PPT one has to keep track which operators can be built up by products of $\tilde{A}_i, \tilde{B}_j$ and linear combinations of them, or, more precisely, by the operators set

$$\mathcal{Q} = \left\{ Q \in \mathbb{B}(\tilde{\mathcal{H}}) \middle| Q = \sum_{ij} c_{ij} \tilde{A}(i)\tilde{B}(j), c_{ij} \in \mathbb{C} \right\}. \tag{22}$$

From the given finite-dimensional commuting Hermitian operators $\tilde{A}_i, \tilde{B}_j$ one can infer the following decompositions

$$\begin{aligned}
\tilde{\mathcal{H}} &= \oplus_k \mathcal{H}_{AB,k} \otimes \mathcal{K}_k &(23)\\
&= \oplus_k (\oplus_l \mathcal{H}_{A,kl} \otimes \mathcal{H}_{B,kl}) \otimes \mathcal{K}_k, &(24)
\end{aligned}$$

and representations

$$\begin{aligned}
\mathcal{Q} &= \oplus_k \mathbb{B}(\mathcal{H}_{AB,k}) \otimes \mathbb{1}, &(25)\\
\tilde{A}_i &= \oplus_k A_{i,k} \otimes \mathbb{1} = \oplus_k (\oplus_l \tilde{A}_{i,kl} \otimes \mathbb{1}) \otimes \mathbb{1}, &(26)\\
\tilde{B}_j &= \oplus_k B_{j,k} \otimes \mathbb{1} = \oplus_k (\oplus_l \mathbb{1} \otimes \tilde{B}_{j,kl}) \otimes \mathbb{1}. &(27)
\end{aligned}$$

These statements can be inferred for instance from Theorem A.7 of Ref. [31]: The original structure given by Eq. (23) comes from the decomposition of (the finite-dimensional C^*-algebra) $\mathcal{Q}$. The tensor product structure of $\mathcal{H}_{AB,k}$, included in Eq. (24), originates from the same argument as in the standard Tsirelson problem, applied to the operators $\tilde{A}_{i,k}, \tilde{B}_{j,k}$ acting on $\mathcal{H}_{AB,k}$ which still commute. (In addition to Theorem A.7 of Ref. [31] one needs to know that $\tilde{B}_{j,k}$ are elements of the commutant of C^*-algebra generated by $\{\tilde{A}_{i,k}\}$).

Since it will become important shortly, let us stress the meaning of Eq. (25): It states that any operator of this form is in the set $\mathcal{Q}$. Note that this structure includes, in particular, all operators $\oplus_k[\oplus_l Q_{kl}] \otimes \mathbb{1}$ with Q_{kl} arbitrary. Thus, also the very special operators where all of these operators Q_{kl} are zero except for one particular pair, maybe $k = k', l = l'$. Hence by appropriately chosen coefficients $c_{ij}[Q_{k'l'}]$ it is possible to build up

$$\sum_{ij} c_{ij}[Q_{k'l'}] \tilde{A}(i)\tilde{B}(j) = Q_{k'l'} \otimes \mathbb{1}_{\mathcal{K}_{k'}} \tag{28}$$

with $Q_{k'l'}$ being an arbitrary operator acting on $\mathcal{H}_{AB,kl}$.

Next let us construct the bipartite state. In the following we denote by Π_{kl} the projections onto $\mathcal{H}_{AB,kl} \otimes \mathcal{K}_k$.

From the structure of the Hilbert space and the measurements one obtains the identity

$$\mathrm{tr}[\tilde{\rho}\tilde{A}_i\tilde{B}_j]=\sum_{kl}\mathrm{tr}_{AB,kl}[\mathrm{tr}_{\mathcal{K}_k}(\Pi_{kl}\tilde{\rho}\Pi_{kl})\tilde{A}_{i,kl}\otimes\tilde{B}_{j,kl}]. \quad (29)$$

If one defines $p_{kl}\tilde{\rho}_{AB,kl} = \mathrm{tr}_{\mathcal{K}_k}(\Pi_{kl}\tilde{\rho}\Pi_{kl})$ one can interpret Eq. (29) as a classical mixture of bipartite states on different subsystems. Via appropriate local auxiliary system one can combine this into a single bipartite system

$$\rho_{AB} = \sum p_{kl}\tilde{\rho}_{AB,kl} \otimes |kl,kl\rangle_{A'B'}\langle kl,kl|\,, \quad (30)$$

$$A_i = \sum \tilde{A}_{i,kl} \otimes |kl\rangle_{A'}\langle kl|\,, \quad (31)$$

$$B_j = \sum \tilde{B}_{j,kl} \otimes |kl\rangle_{B'}\langle kl|\,, \quad (32)$$

such that $\mathrm{tr}[\tilde{\rho}\tilde{A}_i\tilde{B}_j] = \mathrm{tr}[\rho_{AB}A_i \otimes B_j]$ following Eq. (29). That ρ_{AB} is indeed a valid quantum state, while A_i, B_j are correct operators describing measurements follows from their structure.

Thus we are left to show that ρ_{AB} is PPT, which, by its form given by Eq. (30), is equivalent to $\tilde{\rho}_{AB,kl}$ being PPT for all k,l. A finite-dimensional operator $\tilde{\rho}^{T_A}_{AB,kl} \geq 0$ is positive semidefinite if and only if for all operators $Q_{kl} \in \mathbb{B}(\mathcal{H}_{AB,kl})$ it holds that $\mathrm{tr}[\tilde{\rho}^{T_A}_{AB,kl}Q_{kl}Q^\dagger_{kl}] \geq 0$ [38]. In the remaining we show that this inequality holds due to $\mathrm{tr}[\tilde{\rho}QQ^\dagger] \geq 0$ for all operators $Q \in \mathcal{Q}$ and the extra conditions given by Eq. (20). Here it is important to stress that $\mathcal{Q}$ indeed includes all operators $Q_{kl} \in \mathbb{B}(\mathcal{H}_{AB,kl})$ as remarked previously. Using $Q \in \mathcal{Q}$ with $c_{ij} = c_{ij}[Q_{k'l'}]$ as given in Eq. (28) for fixed k', l', $\tilde{A}_{kl}(i) = \tilde{A}_{i_1,kl}\ldots\tilde{A}_{i_n,kl}$ and a similar shorthand for $\tilde{B}_{kl}(j)$, one obtains

$$\begin{aligned}
\mathrm{tr}[\tilde{\rho}QQ^\dagger] &= \sum c_{ij}c^*_{uv}\,\mathrm{tr}[\tilde{\rho}\tilde{A}(i)\tilde{B}(j)(\tilde{A}(u)\tilde{B}(v))^\dagger] \\
&= \sum c_{ij}c^*_{uv}\,\mathrm{tr}[\tilde{\rho}\tilde{A}(i)\tilde{A}(u^T)\tilde{B}(j)\tilde{B}(v)^\dagger] \\
&\overset{\text{Eq. (20)}}{=} \sum c_{ij}c^*_{uv}\,\mathrm{tr}[\tilde{\rho}\tilde{A}(u)\tilde{A}(i^T)\tilde{B}(j)\tilde{B}(v)^\dagger] \\
&= \sum c_{ij}c^*_{uv}\,\mathrm{tr}[\tilde{\rho}(\tilde{A}(i)\tilde{A}(u)^\dagger)^T\tilde{B}(j)\tilde{B}(v)^\dagger] \\
&= \sum c_{ij}c^*_{uv}p_{kl} \\
&\quad \times \mathrm{tr}_{AB,kl}[\tilde{\rho}_{AB,kl}(\tilde{A}_{kl}(i)\tilde{A}_{kl}(u)^\dagger)^T \otimes \tilde{B}_{kl}(j)\tilde{B}_{kl}(v)^\dagger] \\
&= \sum c_{ij}c^*_{uv}p_{kl} \\
&\quad \times \mathrm{tr}_{AB,kl}[\tilde{\rho}^{T_A}_{AB,kl}\tilde{A}_{kl}(i)\tilde{A}_{kl}(u)^\dagger \otimes \tilde{B}_{kl}(j)\tilde{B}_{kl}(v)^\dagger] \\
&= p_{k'l'}\,\mathrm{tr}_{AB,k'l'}[\tilde{\rho}^{T_A}_{AB,k'l'}Q_{k'l'}Q^\dagger_{k'l'}] \geq 0. \quad (33)
\end{aligned}$$

Here we employ the following: Because $\tilde{A}_i$ commutes with every $\tilde{B}_j$ this property naturally extends to products $[\tilde{A}(i), \tilde{B}(j)] = 0$. Moreover, since the measurement operators $\tilde{A}_i$ can be taken real and symmetric, it follows that $\tilde{A}(i)^\dagger = \tilde{A}(i^T)$ and similarly $\tilde{A}(k)\tilde{A}(i^T) = [\tilde{A}(i)\tilde{A}(k)^\dagger]^T$. As indicated the identity given by Eq. (20) is employed in the third line. Afterwards one uses once more the block decomposition as already shown previously, and the identity $\mathrm{tr}[XY^{T_A}] = \mathrm{tr}[X^{T_A}Y]$. The final inequality stems from the positivity of the original density operator $\tilde{\rho}$. This finishes the proof of Proposition 2. □

Let us now see how the property of a PPT mixture can be cast into a commutative version. The equivalence of both formulations in the finite case is shown similarly as in the bipartite case.

Definition 2. *i) An n-partite probability distribution P has a* tensor product PPT mixture quantum representation *if there exists a real n-partite state ρ, local real measurement operators $A^1_i, A^2_j, \ldots, A^n_k$ such that $P = \mathrm{tr}[\rho A^1_i \otimes A^2_j \otimes \ldots \otimes A^n_k]$ and $\rho = \sum_m p_m\rho_m$ being a convex combination of real, PPT invariant states $\rho_m = \rho_m^{T_m}$ for all bipartitions m.*

ii) An n-partite probability distribution P has a commutative PPT mixture quantum representation *if there exists a real state $\tilde{\rho}$, commuting real measurement operators $\tilde{A}^1_i, \tilde{A}^2_j, \ldots, \tilde{A}^n_k$ such that $P = \mathrm{tr}[\tilde{\rho}\tilde{A}^1_i\tilde{A}^2_j\ldots\tilde{A}^n_k]$ and $\rho = \sum_m p_m\rho_m$ is a convex combination of real "PPT states" ρ_m for all possible formal bipartitions m, i.e., each state ρ_m must satisfy that for all possible products of operators from partition m, indexed by $i^{(m)} = (i^{k_1}_1, i^{k_2}_2, \ldots, i^{k_r}_r)$ with $k_s \in m$ for $s = 1, \ldots, r$, $\tilde{A}(i^{(m)}) = \tilde{A}^{k_1}_{i_1}\tilde{A}^{k_2}_{i_2}\ldots\tilde{A}^{k_r}_{i_r}$ and similar for products for its complement m_c it holds*

$$\mathrm{tr}[\tilde{\rho}_m\tilde{A}(i^{(m)})\tilde{A}(j^{(m_c)})] = \mathrm{tr}[\tilde{\rho}_m\tilde{A}(i^{(m)T})\tilde{A}(j^{(m_c)})] \quad (34)$$

with $i^{(m)T} = (i^{k_r}_r, i^{k_{r-1}}_{r-1}, \ldots, i^{k_1}_1)$.

Proposition 3. *Any probability distribution which has a tensor product PPT mixture quantum representation also has a commutative PPT mixture quantum representation. The converse holds at least if the commutative PPT mixture quantum representation is finite-dimensional.*

Proof. The direction from *i*) to *ii*) holds by similar arguments as given in the bipartite case.

For the converse direction one proceeds analogously as in the proof of Proposition 2. From the given measurement operators one can obtain, similar to Eqs.(23)-(27), a decomposition of the underlying Hilbert space given by

$$\mathcal{H} = \oplus_k(\oplus_l\mathcal{H}_{A_1,kl} \otimes \mathcal{H}_{A_2,kl} \otimes \ldots \otimes \mathcal{H}_{A_n,kl}) \otimes \mathcal{K}_k, \quad (35)$$

its measurement operators and respective operator set $\mathcal{Q}$, i.e., the set of all operators generated by linear combinations of products of the measurements [31]. This shows that all measurements $\tilde{A}^s_i$ from system s act only nontrivial on the part $\mathcal{H}_{A_s,kl}$, in analogy to the bipartite case as given by Eqs. (26),(27). Via the projector Π_{kl} onto the subspace $\mathcal{H}_{A_1\ldots A_n,kl} \otimes \mathcal{K}_k$ one arrives again at the tensor product form

$$\begin{aligned}
&\mathrm{tr}[\tilde{\rho}\tilde{A}^1_i\ldots\tilde{A}^n_j] \\
&\quad = \sum_{kl} p_{kl}\,\mathrm{tr}_{A_1\ldots A_n,kl}[\tilde{\sigma}_{kl}\tilde{A}^1_{i,kl} \otimes \ldots \otimes A^n_{j,kl}]. \quad (36)
\end{aligned}$$

with $p_{kl}\tilde{\sigma}_{kl} = p_{kl}\tilde{\sigma}_{A_1\ldots A_n,kl} = \mathrm{tr}_{\mathcal{K}_k}[\Pi_{kl}\tilde{\rho}\Pi_{kl}]$. Via auxiliary states this classical mixture can be turned into a single multipartite state, i.e., $\rho = \sum_{kl} p_{kl}\tilde{\sigma}_{kl} v \otimes |kl\rangle\langle kl|^{\otimes n}$ and measurements $A_i^s = \sum_v \tilde{A}_{i,kl}^s \otimes |kl\rangle\langle kl|$ for each system s.

At last one needs to show that ρ is a PPT mixture, which amounts to verifying that each $\tilde{\sigma}_{kl}$ is a PPT mixture. This decomposition is given by the original expansion $\tilde{\rho} = \sum_m \tilde{\rho}_m$ into real (in this case un-normalized) states $\tilde{\rho}_m$ which satisfy the property given by Eq. (34). If we set $\rho_{m,kl} = \mathrm{tr}_{\mathcal{K}_k}[\Pi_{kl}\tilde{\rho}_m\Pi_{kl}]$ then one obtains a decomposition

$$p_{kl}\tilde{\sigma}_{kl} = \sum_m \tilde{\rho}_{m,kl}, \tag{37}$$

which verifies that $\tilde{\sigma}_{kl}$ is a PPT mixture if one can show that $\tilde{\rho}_{m,kl}$ is PPT with respect to bipartition m. Since we can interpret $\tilde{\rho}_{m,kl}$ as a bipartite system on m and its complement $\bar{m}$ this boils down to the bipartite argument: The condition $\tilde{\rho}_{m,kl}^{T_m} \geq 0$ is equivalent to $\mathrm{tr}[\tilde{\rho}_{m,kl}^{T_m} Q_{kl} Q_{kl}^\dagger] \geq 0$ for all operators $Q_{kl} \in \mathbb{B}(\mathcal{H}_{A_1\ldots A_n,kl})$. Since any operator Q_{kl} can again be written as a linear combination of products of the measurement operators, the identities given by Eq. (34) verify $\mathrm{tr}[\tilde{\rho}_{m,kl}^{T_m} Q_{kl} Q_{kl}^\dagger] = \mathrm{tr}[\tilde{\rho}_{m,kl} Q_{kl} Q_{kl}^\dagger] \geq 0$, where the inequality results from the positivity of $\tilde{\rho}_m$. This concludes the proof. ☐

Finally let us state the convergence result. As already pointed out this is a corollary of the original convergence proof for the NPA hierarchy [41]. In this part we use the notation $\chi^{(\ell)}$ to refer to the matrix of moments of level ℓ, i.e., having for each party a product of up to ℓ local operators in the construction. We first prove the convergence of the PPT hierarchy for the bipartite case. This proof then directly generalizes to a PPT hierarchy for a single bipartition m in the multipartite case, which finally helps to prove the convergence of the PPT mixture in the end. Note that the condition $\chi^{(\ell)}[P,u]_{\mathrm{tr}} = 1$ does not appear since we explicitly talk about probability distributions.

Proposition 4. *If a bipartite distribution P satisfies $\chi^{(\ell)}[P,u] = \chi^{(\ell)}[P,u]^{T_{\bar{A}}} = \chi^{(\ell)}[P,u]^T \geq 0$ via suitable u for all levels ℓ, then it has a (potentially infinite-dimensional) commutative PPT quantum representation.*

Proof. In the following we use the label s, t, v, w to denote sequences of measurement operators, e.g., $s = (s_1, s_2, \ldots, s_n)$ with $A(s) = A_{s_1}A_{s_2}\ldots A_{s_n}$. However, if we employ i, j it refers to a single index, and $i = 0$ or $j = 0$ should correspond to the identity operator. The structure of $\chi^{(\ell)}$, as given by Eq. (1) for $\ell = 1$, is such that the matrix elements correspond to expectation values

$$\langle st|\chi^{(\ell)}|vw\rangle = \mathrm{tr}\{\rho[A(s)\otimes B(t)][A(v)\otimes B(w)]^\dagger\}. \tag{38}$$

The identities given by the described partial information can be parsed as $\langle st|\chi^{(\ell)}|vw\rangle = \langle s't'|\chi^{(\ell)}|v'w'\rangle$ if $A(s)A(v)^\dagger \otimes B(t)B(w)^\dagger = A(s')A(v')^\dagger \otimes B(t')B(w')^\dagger$ by the properties satisfied by operators, which are listed as 1) – 3) in the main text. The two extra requirements provide the additional relations $\langle st|\chi^{(\ell)}|vw\rangle = \langle vt|\chi^{(\ell)}|sw\rangle = \langle vw|\chi^{(\ell)}|st\rangle$.

Next let us summarize the convergence idea of the NPA hierarchy [41]. Suppose that $\chi^{(\ell)}[P,u] = \chi^{(\ell)}[P,u]^{T_{\bar{A}}} = \chi^{(\ell)}[P,u]^T \geq 0$ exists for each level ℓ, then there is also a limit $\chi^{(\infty)} \geq 0$ which satisfies all the listed identities. For this infinite-dimensional matrix one can associate a set of vectors $\{|e_{st}\rangle\}$ with $\langle st|\chi^{(\infty)}|vw\rangle = \langle e_{st}|e_{vw}\rangle$. Via this set one now defines the measurements and the corresponding state. The measurements are $\hat{A}_i = \mathrm{proj}(\{|e_{st}\rangle : s_1 = i\})$ where proj stands for the projector onto the corresponding set. Similarly one defines $\hat{B}_j = \mathrm{proj}(\{|e_{st}\rangle : t_1 = j\})$, while the state is given by $\hat{\rho} = |e_{00}\rangle\langle e_{00}|$. As shown in detail in Ref. [41] these measurements satisfy $\hat{A}_i|e_{st}\rangle = |e_{\tilde{s}t}\rangle$ with $\tilde{s} = (i, s)$ and similar for $\hat{B}_j$ due to the linear constraints. These choices reproduce the observed expectation values since

$$\begin{aligned} P_{ij} &= \langle e_{00}|e_{ij}\rangle = \langle e_{00}|\hat{A}_i\hat{B}_j|e_{00}\rangle \\ &= \mathrm{tr}[|e_{00}\rangle\langle e_{00}|\hat{A}_i\hat{B}_j] = \mathrm{tr}[\hat{\rho}\hat{A}_i\hat{B}_j]. \end{aligned} \tag{39}$$

Moreover, besides being true projectors, the operators $\hat{A}_i, \hat{B}_j$ fulfil the projector identity and commute.

Now let us add the remaining properties. First, let us point out that since $\chi^{(\infty)} = \chi^{(\infty)T}$ the set $\{|e_{st}\rangle\}$ can be chosen to be a set of real vectors. Therefore, the state and also the measurements are real symmetric, which implies in particular $\hat{A}(s)^\dagger = \hat{A}(s^T)$. Thus we are left to show the property given by Eq. (20). However, this follows from the extra requirement $\langle st|\chi^{(\infty)}|vw\rangle = \langle vt|\chi^{(\infty)}|sw\rangle$ since

$$\begin{aligned} \mathrm{tr}[\hat{\rho}\hat{A}(s)\hat{B}(t)] &= \langle e_{00}|\hat{A}(s)\hat{B}(t)|e_{00}\rangle \\ &= \langle e_{00}|e_{st}\rangle = \langle e_{s0}|e_{0t}\rangle = \langle e_{00}|\hat{A}(s)^\dagger\hat{B}(t)|e_{00}\rangle \\ &= \langle e_{00}|\hat{A}(s^T)\hat{B}(t)|e_{00}\rangle = \mathrm{tr}[\hat{\rho}\hat{A}(s^T)\hat{B}(t)]. \end{aligned} \tag{40}$$

This concludes the convergence proof. ☐

Let us remark that the proof works equivalently in the multipartite case if one is only interested in a single bipartition m: If an n-partite distribution P satisfies $\chi^{(\ell)}[P,u] = \chi^{(\ell)}[P,u]^{T_{\bar{m}}} = \chi^{(\ell)}[P,u]^T \geq 0$ via suitable u for all levels ℓ, then it has a (potentially infinite-dimensional) commutative PPT quantum representation with respect to partition m. As for the bipartite case one uses the convergence statement in the multipartite case to show existence of a respective state $\hat{\rho}$ and commuting measurements $\hat{A}_i^1, \ldots, \hat{A}_k^n$. The additional linear requirements given by Eq. (34) for m are shown similarly as in Eq. (40) by using the identities $\chi^{(\ell)}[P,u] = \chi^{(\ell)}[P,u]^{T_{\bar{m}}} = \chi^{(\ell)}[P,u]^T$.

Before we continue we like to clarify another point: As seen in the proof of Proposition 4 one always obtains a pure state $\tilde{\rho} = |\psi\rangle \langle\psi|$ in the commutative version. However, since any pure PPT state of a tensor product Hilbert space is separable and thus possesses a LHV model, one could ask whether this implies that any (finite-dimensional) PPT state must necessarily satisfy all Bell inequalities. This would prove the Peres conjecture, at least, in the finite-dimensional case. However this is not true. Indeed, even if the state in the commutative version is pure, the related bipartite, tensor product state following the procedure outlined in the proof of Proposition 2 does not have to be. More precisely, this bipartite state ρ_{AB} is given by Eq. (30), where the partial trace over $\mathcal{K}_k$ in $\rho_{AB,kl} = \mathrm{tr}_{\mathcal{K}_k}(\Pi_{kl} |\psi\rangle \langle\psi| \Pi_{kl})$ results in a mixed PPT state. This extra system $\mathcal{K}_k$ can be considered as a purifying system of each bipartite state $\rho_{AB,kl}$ to $\Pi_{kl} |\psi\rangle$, which would then be pure if one translates it into commutative terms. Finally note that the purifying systems $\mathcal{K}_k$ would vanish if the commuting measurements $\tilde{A}_i$, $\tilde{B}_j$ would generate $\mathbb{B}(\tilde{\mathcal{H}})$.

At last, we prove the convergence of the characterization of an n-partite distribution P with an underlying PPT mixture, which shows that the hierarchy given by Eq. (8) is complete.

Proposition 5. *If an n-partite distribution P satisfies $\chi^{(\ell)}[P,u] = \sum \chi^{(\ell)}[P_m, u_m]$ with $\chi^{(\ell)}[P_m, u_m] = \chi^{(\ell)}[P_m, u_m]^{T_{\bar{m}}} = \chi^{(\ell)}[P_m, u_m]^{T} \geq 0$ via suitable P_m, u_m for all levels ℓ, then it has a (potentially infinite-dimensional) commutative PPT mixture quantum representation.*

Proof. From the remark after Proposition 4 we know that the conditions on $\chi^{(\ell)}[P_m, u_m]$ for all ℓ show that there exists an (un-normalized) distribution P_m which has a commutative PPT quantum representation for the bipartition m. Let us refer to this representation as the state σ_m and measurements $A_i^{1|m}, A_j^{2|m}, \ldots, A_k^{n|m}$. Note that here, because the probabilities P_m are not normalized but we rather have $\sum_m P_m = P$, the state ρ_m satisfies $\mathrm{tr}(\rho_m) = \chi[P_m, u_m]_{\mathrm{tr}}$.

From this one can construct a commutative PPT mixture representation for the given n-partite distribution P by using $\mathcal{H} = \oplus_m \mathcal{H}_m$, $\rho = \oplus_m \sigma_m$ and $A_i^s = \oplus_m A_i^{s|m}$ for each system s. This construction directly ensures positivity and that the measurements commute. A commutative PPT mixture decomposition of the state is then given by $\rho = \sum_m \rho_m$ with (un-normalized) states $\rho_m = \Pi_m \rho \Pi_m$ and Π_m denoting the projector onto $\mathcal{H}_m$. That these states ρ_m indeed satisfy the relations given by Eq. (34) follows from the corresponding identities on σ_m since $\rho_m A_i^s = \sigma_m A_i^{s|m}$ for each system s and similarly for products of more operators. This concludes the proof. $\square$